



POSTER PRESENTATION

Open Access

# Phenotypic variability in a cohort of 40 Italian subjects carrying mutations in the gene *EDA*

L Guazzarotti<sup>1\*</sup>, G Tadini<sup>2</sup>, GE Mancini<sup>3</sup>, S Giglio<sup>4</sup>, I Sani<sup>4</sup>, P Nannini<sup>1</sup>, M Bosoni<sup>1</sup>, A Bottero<sup>5</sup>, A Caimi<sup>5</sup>, M Morelli<sup>1</sup>, GV Zuccotti<sup>1</sup>

From 5th International Conference on Ectodermal Dysplasia (ED2012)  
Erlangen, Germany. 1-3 June 2012

Ectodermal dysplasias (ED) are a group of clinically and genetically heterogeneous conditions commonly characterized by abnormal development of at least two structures derived from the embryonal ectoderm (hair, teeth, nails, and sweat glands). X-linked hypohidrotic ED, which is caused by mutations in the gene *EDA* (MIM 305100), is the most frequent form. In this study, we investigated the phenotype of 40 male patients, aged 2 to 20 years, who all showed developmental defects of ectodermal derivatives and a mutation in *EDA*. Specialist assessments of the involved organ systems were performed. 95% of these patients presented with impairment of sweating, which was only moderate in 35%, while two subjects did not show any alteration of sweat gland function. Severe oligodontia was found in 80% of the subjects, 10% had hypodontia. 90% of the patients showed abnormal crown morphology of the teeth. Severe involvement of the scalp hair was observed in 22% of the patients, moderate involvement in 72%, and no relevant alterations of hair morphology, quantity or growth in 2 patients. Onychodystrophy was seen in 67% of our patients. Concerning minor alterations of ectodermal tissues, we found dry eye signs in 92% of the subjects investigated, recurrent respiratory infections in 82%, hearing loss in 10%, atopic dermatitis in 35%, and a neuropsychological disorder in 10%. Our study shows that an *EDA* mutation can be present in males also in the context of only one of the major clinical signs of X-linked ED, but associated with minor alterations. Therefore we suggest that *EDA* gene analysis should be considered also for males with a mild ED phenotype.

#### Author details

<sup>1</sup>Department of Pediatrics, University of Milan, Italy. <sup>2</sup>IRCCS Ca' Granda Policlinico Hospital, Milan, Italy. <sup>3</sup>IRCCS Galeazzi, Milan, Italy. <sup>4</sup>University of Florence, Italy. <sup>5</sup>University of Milan, Italy.

Published: 25 May 2012

doi:10.1186/1746-160X-8-S1-P2

**Cite this article as:** Guazzarotti et al.: Phenotypic variability in a cohort of 40 Italian subjects carrying mutations in the gene *EDA*. *Head & Face Medicine* 2012 **8**(Suppl 1):P2.

**Submit your next manuscript to BioMed Central and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
[www.biomedcentral.com/submit](http://www.biomedcentral.com/submit)



<sup>1</sup>Department of Pediatrics, University of Milan, Italy  
Full list of author information is available at the end of the article