



RESEARCH

Open Access

Psychological profile and self-administered relaxation in patients with craniofacial pain: a prospective in-office study

Christian Kirschneck^{1*}, Piero Römer¹, Peter Proff¹ and Carsten Lippold²

Abstract

Introduction: The objective of this study was to evaluate the psychological profile of craniofacial pain sufferers and the impact of patient subtype classification on the short-time effectiveness of a self-administered relaxation training.

Methods: One hundred unselected in-office patients (67% females) suffering from chronic facial pain and/or headache with the presumptive diagnose of temporo-mandibular disorder (TMD) completed a questionnaire battery comprising craniofacial pain perception, somatic complaints, irrational beliefs, and pain behavior and were classified into subtypes using cluster analysis. They underwent a self-administered progressive relaxation training and were re-evaluated for pain perception after 3 months.

Results: Pain was mild to moderate in the majority of patients. Symptom domains comprised parafunctional activities, temporo-mandibular pain and dysfunction, fronto-temporal headache, head/neck and neck/back pain. Three patient subtypes were identified regarding symptom/dysfunction level: (i) low burden (mild/moderate), (ii) psychosocial dysfunction (moderate/high), (iii) adaptive coping (moderate/mild). Self-rated adherence to the recommended relaxation training was moderate throughout the sample, but self-rated relief was significantly different between clusters. At follow-up, pain intensity was significantly decreased in all patients, whereas pain-related interference was improved only in dysfunctional and adaptive patients. Improvement of symptom domains varied between clusters and was most comprehensive in adaptive patients.

Conclusions: In conclusion, craniofacial pain sufferers can be divided in meaningful subtypes based on their pain perception, irrational beliefs, and pain behaviour. A self-administered relaxation training generally yielded positive effects on pain perception, however the benefit may be greater in patients with more marked symptom impact (both dysfunctional and adaptive).

Keywords: Craniofacial pain, Temporomandibular dysfunction, Patient classification, Relaxation training, Psychological features

Introduction

Orofacial pain is a common problem and frequently related to temporomandibular dysfunction (TMD) or generalized musculoskeletal pain [1-4]. Head-neck pain distinct from that in the masticatory muscles is among the leading accessory signs in patients with myogenic cranio-mandibular dysfunctions [5-8]. The causes of these disorders are poorly understood, however female

gender and psychosocial dysfunction have been consistently reported as significant risk factors in the development and maintenance of persistent craniofacial pain [5,9-13]. Many pain problems presented to the dentist respond to treatment or resolve on their own. However, pain and disability significantly influenced by psychosocial factors are unlikely to improve substantially with conventional dental treatment alone [14,15]. When psychosocial factors are involved but not recognized, craniofacial pain may become chronic and set up a revolving door, with the patient visiting numerous providers in the wake of treatment failures that may even exacerbate patient distress

* Correspondence: christian.kirschneck@ukr.de

¹Department of Orthodontics, University Medical Centre of Regensburg, Franz-Josef-Strauß-Allee 11, Regensburg 93053, Germany
Full list of author information is available at the end of the article

[16]. Thus, the well-supported bio-psychosocial perspective on chronic pain has been accepted widely [14,17-19] and should be integrated into the dental curriculum [20]. According to this model, a patient's experience of pain and treatment responses are affected not only by nociceptive processes, but also by factors such as mood, beliefs and appraisals as well as by social responses of others. Since, therefore, chronic pain sufferers are not a psychologically homogeneous group [19,21-24], treatment outcome depends upon patient subtype profiles [25-30].

A wide range of therapeutic approaches to craniofacial pain has been proposed, including physical therapies, drugs, and bio-behavioral modalities [31-33]. Even though a dentist is seen by the majority of patients, many of them receive additional non-dental treatment such as relaxation training [31,32,34]. The rationale is that pain and oral parafunctions may be related to stress, poor muscle discrimination or unconscious bracing of the orofacial muscles [35]. Derived from behavioral psychology, relaxation treatment aims to break existing muscle tension pain cycles and, in addition, to provide the patient with a method for controlling pain [36]. Thus, relaxation may be used as a coping strategy to be applied in everyday situations rather than in a treatment setting. Consequently, relaxation is among the most commonly used self-care techniques in chronic pain sufferers with myofascial TMD [37,38].

The present study aims to classify unselected in-office patients with persistent craniofacial pain for their self-perceived pain experiences and psychological profiles (irrational attitudes, coping behavior) and to evaluate the short-term outcomes of a self-administered relaxation training in the patient subtypes identified.

Materials and methods

Patients

The study group comprised 100 craniofacial pain sufferers between 20 and 75 years of age with a suspected diagnosis of TMD. Volunteers from general dental offices who gave their informed consent were randomly entered into the study in the order of their appointments. In order to ensure patient anonymity, questionnaire data were not linked to clinical records. The research was conducted in accordance with the declaration of Helsinki and the ethical regulations of the University of Regensburg, Germany.

Questionnaires

The questionnaire battery comprised the components shown in Table 1.

The Pain Inventory (PI) involves items that were adopted from an existing clinical ad-hoc questionnaire to assess the type and extent of cranial and facial pain during the last fortnight. General pain ratings (PI-A1 to -A4)

were obtained for current intensity, frequency (per week), duration (hours per day) and pain-related interference. The main questionnaire body included of a total 28 items on five factor-analytic derived subscales (PI-1 to PI-5) with sufficient internal consistency (Cronbach's α ranging from 0.77 to 0.87).

The German version [39] of the McGill Pain Questionnaire MPQ was used to assess the affective (MPQ-A) and sensory (MPQ-S) dimensions of pain perception.

Zerssen's Complaint List (CL) is a well-proven German self-assessment instrument involving 24 somatic complaints such as nausea, shortness of breath, muscle pain, and psychosomatic symptoms such as irritability, sleep disorder and uneasiness [40].

The Irrational Attitudes Questionnaire comprises a total of 30 items on 4 subscales with good reliability and validity [41].

The Pain Behavior Questionnaire [42] includes 29 items related to pain and coping behaviors classified into 4 subscales with internal consistencies ranging between $\alpha = 0.68$ and 0.84 in rheumatic and headache patients.

Treatment

Patients were provided with an audio tape containing instructions for progressive muscle relaxation after Edmund Jacobson. The following areas of systematic tensing and relaxing of muscle groups were included in subsequent order:

1. both hands and arms,
2. face (including the jaw muscles), neck and shoulders
3. chest, back and belly
4. both thighs, shanks and feet.

The participants were advised to listen to and follow the exercises as often as possible. After 3 months, a follow-up assessment was carried out including the Pain Inventory and the McGill Pain Questionnaire to allow for evaluation of the treatment effects. Moreover, patient satisfaction and compliance with the relaxation training as well as pain-relief was assessed using graded rating scales.

Statistical analysis

Statistical data analysis using SPSS for Windows software included descriptive statistics of variable distributions, Ward's cluster analysis of subjects followed by k-means procedures, and significance testing for between-group (Kruskal-Wallis H test) and pre-post-treatment effects (Wilcoxon signed ranks test). The general significance level was set to $\alpha < 0.05$.

Table 1 Questionnaires used in the study

Items/Subscales	Items	rating scale
Pain inventory		
<i>General items:</i>		
PI-A1	Item: <i>intensity of pain</i>	1 (not noticeable) – 6 (unbearable)
PI-A2	Item: <i>frequency of pain</i>	1 (daily) – 4 (once in a fortnight)
PI-A3	Item: <i>duration of pain</i>	1 (less than 1 hour) – 7 (permanently)
PI-A4	Item: <i>pain-related interference</i>	1 (unrestricted) – 5 (no activities)
<i>Subscales:</i>		
PI-1	Subscale: <i>"parafunctional activities (PFA)"</i>	1 (not at all) – 4 (very much)
PI-2	Subscale: <i>"temporomandibular dysfunction (TMD)"</i> chewing difficulties, myofascial pain, temporomandibular pain and stiffness	7
PI-3	Subscale: <i>"cervicodorsal pain (CDP)"</i>	8
PI-4	Subscale: <i>"occipitocervical pain (OCP)"</i> including TMJ sounds	5
PI-5	Subscale: <i>"frontotemporal pain (FTP)"</i> including migraine-like headache	5
McGill pain questionnaire		
MPQ-A	subscale: <i>affective</i> descriptors	7 (not at all) – 4 (very severely)
MPQ-S	subscale: <i>sensory</i> descriptors	14
Zerssen complaint list		
		10
Irrational attitudes questionnaire		
		24 (not at all) – 4 (severely)
IA-1	<i>"negative self-appraisal"</i> (of own person and abilities)	0 (strongly disagree) – 5 (completely agree)
IA-2	<i>"dependency"</i> (of own behavior upon approval by others)	8
IA-3	<i>"internal attribution"</i> (of problems to own responsibility)	8
IA-4	<i>"irritability"</i> (due to external stressors)	7
Pain behavior questionnaire		
		7 (strongly disagree) – 5 (completely agree)
PB-1	<i>"avoidance"</i> (of social contact, leisure and work activities)	8
PB-2	<i>"cognitive control"</i> (of pain using relaxation, attention and mind techniques)	8
PB-3	<i>"social support"</i> (perceived by patient)	6
PB-4	<i>"activity"</i> (in social and professional areas)	7

Results

Pretreatment findings

General patient features

Cluster analysis involving all questionnaire scales yielded a classification of 98% of the interviewees into three significant subgroups comprising 27 (I), 28 (II) and 43 (III) subjects respectively.

Sixty-six interviewees were females, thirty-two were males, while gender was not specified in two cases. The female-to-male ratio was 1.5 in cluster I, 3.7 in cluster II and 1.9 in cluster III, yet not significantly different. Most of the interviewees were married (47%) or coupled (18%), while 27% were single and 5% were divorced.

Patient age ranged from 20 to 75 years (mean \pm standard deviation 37.7 ± 13.7 years), the interquartile range was from 28 to 46 (median 32 years). The duration of illness

was 6.7 ± 6.9 years on average. The pain symptoms had existed for 6 months to 20 years (median 5 years) in 90% of subjects.

Patient age and duration of illness (Figure 1) were significantly different between clusters ($p \leq 0.02$). Mean age was 38.4 years in cluster I, 31.7 years in cluster II, and 40.9 years in cluster III. The pain symptoms had existed for an average of 4.25 (I), 5.5 (II) and 8.9 (III) years respectively.

Pain scales

Pain intensity (PI-A1) was rated as "mild" by 35%, "moderate" by 36% and "severe" by 16% of the subjects. Pain was "not noticeable" or "very severe" in 4% respectively and "intolerable" in 2 cases. Pain intensity was significantly higher in clusters II and III compared to cluster I.




Figure 1 Patient age at assessment (solid line) and at reported onset of symptoms (broken line).

The frequency of pain episodes showed a modal value of “1-3 times a week” (36%), followed by “once in a fortnight” (26%). Daily pain episodes were reported less frequently (14%) by the interviewees, while 19% stated that they experienced pain 4-6 times a week.

The pain episodes lasted for “less than 1 hour” or “about 1-6 hours” in 34% respectively. A longer duration was stated by 28%, *i.e.* “about 6-8 hours” (14%), “about 12-24 hours” (12%) and “continuously” (2%).

Also, interference of pain with daily life activities was assessed. All activities were possible in 77% of the interviewees (“pain noticed when attended to” 24%, “pain temporarily disregarded” 26%, “pain noticed during all activities” 27%). Due to pain intensity, 11% were “restricted to light activities” and 8% were “restricted from any activities”. Pain-related interference was significantly greater in clusters II and III compared to cluster I.

The overall parameters for the Pain Inventory and McGill Pain Questionnaire items broken down by cluster membership are summarized in Table 2.

The overall item medians for symptom type and location were 1.5 (corresponding to “not at all” or “a little”) for TMD and 2.0 (corresponding to “a little”) for the other symptom subscales (PI-1, PI-3 to PI-5), respectively.

Patients of cluster II displayed significantly higher values for TMD (PI-2) and fronto-temporal headache (PI-5) compared to the other clusters. CDP (PI-3) was significantly less marked in cluster I compared to the clusters II and III.

The total item medians for affective (MPQ-A) and sensory (MPQ-S) pain perception were 1.5 (corresponding to

“not at all” and “a little”). Both affective and sensory scores were significantly elevated in cluster II subjects.

Other scales

The overall item median of Zerssen’s Complaint List was 1.5 (corresponding to “not at all” or “scarce”). Scores were significantly higher in clusters II and III compared to cluster I.

Regarding the Irrational Attitudes subscales, the item medians were highest (2.5) for “dependency” (IA-2), lowest (1.5) for “negative self-appraisal” (IA-1) and moderate (2.0) for “internal attribution” (IA-3) and “irritability”

Table 2 Pretreatment findings for pain scales: Md median, IQR interquartile range, p significance of inter-cluster differences (Kruskal-Wallis H-test)

	Cluster I		Cluster II		Cluster III		p	Total	
	Md	IQR	Md	IQR	Md	IQR		Md	IQR
PI-A1	2.0	2.0; 3.0	3.0	2.0; 4.0	3.0	2.0; 3.0	0.02	3.0	2.0; 3.0
PI-A2	3.0	3.0; 4.0	2.0	2.0; 3.0	3.0	2.0; 4.0	ns	3.0	2.0; 4.0
PI-A3	1.0	1.0; 2.0	2.0	2.0; 3.0	2.0	1.0; 3.0	ns	2.0	1.0; 3.0
PI-A4	2.0	1.0; 3.0	3.0	2.0; 4.0	3.0	2.0; 3.0	0.009	2.0	1.5; 3.0
PI-1	2.0	1.5; 2.5	2.0	2.0; 3.0	2.0	1.5; 2.5	ns	2.0	1.5; 2.5
PI-2	1.5	1.0; 2.0	1.5	1.5; 2.0	1.5	1.0; 1.5	0.001	1.5	1.0; 2.0
PI-3	1.5	1.0; 2.0	2.0	2.0; 2.5	2.0	1.5; 3.0	0.002	2.0	1.5; 2.5
PI-4	2.0	1.0; 2.0	2.0	1.5; 2.5	2.0	1.5; 3.0	ns	2.0	1.5; 2.5
PI-5	1.5	1.0; 2.0	2.0	1.5; 2.5	2.0	1.0; 2.0	0.005	2.0	1.5; 2.0
MPQ-A	1.5	1.0; 1.5	2.0	1.5; 2.5	1.5	1.0; 2.0	<0.001	1.5	1.0; 2.0
MPQ-S	1.5	1.0; 1.5	1.5	1.5; 2.0	1.5	1.0; 1.5	0.04	1.5	1.0; 2.0

(IA-4), respectively. All subscale scores were highest in cluster II and lowest in cluster III. These differences were significant.

The item medians of the Pain Behavior subscales were 3.5 for “activity” (PB-4), 3.0 for “cognitive control” (PB-2), and 2.0 for “avoidance” (PB-1) and “social support” (PB-3), respectively. The Pain Behavior scores were not significantly different between clusters except for “avoidance” (PB-1), which was elevated in cluster II subjects.

The overall item parameters for the Complaint List, the Irrational Attitudes Questionnaire and the Pain Behavior Questionnaire are given in Table 3.

Posttreatment findings

Patient-perceived pain was re-evaluated 3 months after instruction for the self-administered relaxation training. While acceptance of (median 4 “much”) and adherence to (median 2 “1-3 times a week”) the program were not significantly different between clusters, self-rated improvement varied between cluster I patients (median 2.5) and cluster II and III patients (median 3.0). Thus, the clusters were significantly ($p = 0.01$) different for self-assessed pain relief after relaxation exercise (Figure 2).

Pre-post analyses of the pain-related scales (H-Test) confirmed global improvements for the total sample, however revealed significant differences between clusters for pain intensity (PI-1 $p = 0.03$), episode duration (PI-3 $p = 0.01$), pain-related interference (PI-4 $p = 0.008$), the affective (MPQ-A $p = 0.04$) and sensory (MPQ-S $p = 0.005$) components of pain perception as well as the item scores of TMD (PI-A2 $p < 0.001$ and FTP (PI-A5 $p < 0.001$).

Pain intensity (PI-A2) significantly decreased in all patient irrespective of cluster membership ($p \leq 0.005$). The frequency (PI-A3) and duration (PI-A4) of pain episodes as well as pain-related interference (PI-A5) were significantly reduced in cluster II and III patients ($p \leq 0.025$),

Table 3 Pretreatment findings for additional scales: Md median, IQR interquartile range, p significance of inter-cluster differences (Kruskal-Wallis H-test)

	Cluster I		Cluster II		Cluster III		p	Total	
	Md	IQR	Md	IQR	Md	IQR		Md	IQR
CL	1.5	1.0; 1.5	2.0	1.5; 2.5	2.0	1.5; 2.0	<0.001	1.5	1.5; 2.0
IA-1	1.5	1.0; 2.5	3.5	2.5; 4.0	1.0	0.5; 1.0	<0.001	1.5	1.0; 2.5
IA-2	3.0	2.5; 3.5	3.5	3.0; 4.5	1.0	1.0; 2.0	<0.001	2.5	1.5; 3.5
IA-3	2.5	2.0; 3.0	3.5	3.0; 4.0	1.0	0.5; 1.5	<0.001	2.0	1.0; 3.0
IA-4	2.5	2.0; 3.0	3.0	2.5; 4.0	1.5	1.0; 1.5	<0.001	2.0	1.5; 3.0
PB-1	2.0	1.5; 2.0	2.5	2.0; 3.5	2.0	1.5; 3.0	0.001	2.0	1.5; 2.5
PB-2	3.0	2.5; 4.0	2.5	2.0; 3.0	2.5	2.0; 3.5	ns	3.0	2.0; 3.5
PB-3	2.0	1.5; 3.0	2.0	1.0; 2.5	2.0	1.5; 3.5	ns	2.0	1.5; 3.0
PB-4	3.5	3.0; 4.0	3.0	2.5; 3.5	3.5	3.0; 4.0	ns	3.5	3.0; 4.0

but not in Cluster I patients. Differential effects were observed for the domains of the Pain Inventory. Cluster I patients showed significant improvements of CDP (PI-3) and FTP (PI-5) and either dimension of the MPQ. In cluster II, significant improvement was achieved for TMD (PI-2) and OCP (PI-4). The most global improvements were revealed for cluster III patients (all scales except TMD). The results are given in Table 4.

Discussion

The multifactorial model of TMD and orofacial pain has prompted numerous studies that emphasize the psychological factors of these disorder and involve alternative treatment approaches [18,31-33]. However, results from dental in-office studies are largely lacking, although most patients have no access to special care institutions.


In order to avoid selection bias, eligible patients with chronic craniofacial pain from general dental practices were consecutively entered into the study and anonymously completed the questionnaire battery. Questionnaires are a well-tried instrument for the assessment of subjective experiences and psychological characteristics in chronic pain [43].

General findings

The sociodemographic characteristics of our sample suggest a 2:1 ratio with females prevailing and a main age range between 28 and 46 years. This distribution corresponds to reports from literature where the highest prevalence of facial pain has been found in females during the main reproductive age from 25 to 40 years [2,5,12,44-46]. Since associated pain areas are also more frequently reported by females [1,47], an increased sensitivity to pain in females has been suggested to account for these findings [10,46].

Almost three thirds of the subjects rated their pain as mild or moderate, while only a minor part reported severe pain and/or daily occurrence of pain which largely prevented them from everyday activities. These findings agree with reports from literature implying that orofacial pain is mild to moderate in 75% of the cases and causes only slight restrictions of everyday activities. Von Korff [48] found more marked restrictions in 16% for facial pain and 30% for headache, while 30-40% of the patients showed the lowest grade of pain intensity. The burden of facial pain assessed in German population surveys corresponds well to our results [44].

Simultaneous occurrence of facial and cranial pain at variable locations has been frequently reported in literature [5-7,49]. Moreover, patients with chronic craniofacial pain frequently show a comorbidity of pain in other regions of the body, especially the neck and back, or other functional complaints which are rarely reported to the dentist [1-4]. Therefore, we utilized Zerssen's CL to



assess the total burden of complaints which was in the abnormal range in more than 90% of the interviewees.

Perceived pain intensity, pain-related interference with daily activities, and behavioral treatment outcome have been shown to be associated with dysfunctional attitudes [50,51]. The frequency of maladaptive, irrational beliefs is increased particularly in patients with facial pain of unknown origin, myofascial pain and daily headache compared to patients with other TMD diagnoses [9,52]. The present unselected sample displayed moderate overall scores which may reflect the relatively low symptom burden and the use of generalized attitudes instead of pain-specific beliefs. Negative “irrational” beliefs are associated with rather passive pain behaviors such as avoidance, inactivity, and social support-seeking [53]. It

is little surprising, hence, that our findings involving pain behavior were generally less dysfunctional as compared to other studies [37,54]. In fact, agreement with statements indicating activity and cognitive control was significantly stronger than with those involving social support-seeking and avoidance.

Treatment response

Self-care and home exercise regimens in orofacial pain sufferers have been encouraged by several studies, all the more so as they are simple and non-invasive, have a favourable cost benefit ratio over other treatment modalities, allow an easy self-management approach and can be managed by the general dental practitioner [14,38,55-57]. However, the provider should not simply hand the patient

Table 4 Posttreatment findings for pain scales: Md median, IQR interquartile range, p significance of pre-posttreatment differences (Wilcoxon signed-rank test)

	Cluster I			Cluster II			Cluster III			Total		
	Md	IQR	p	Md	IQR	p	Md	IQR	p	Md	IQR	p
PI-A1	2.0	2.0; 3.0	0.005	3.0	2.0; 3.5	0.003	2.5	2.0; 3.0	<0.001	2.5	2.0; 3.0	<0.001
PI-A2	3.0	3.0; 4.0	ns	2.5	2.5; 3.0	0.003	3.0	2.5; 4.0	0.005	3.0	2.5; 4.0	<0.001
PI-A3	1.5	1.0; 2.0	ns	2.0	2.0; 3.0	0.008	2.0	1.0; 3.0	0.025	2.0	1.0; 3.0	<0.001
PI-A4	2.0	1.0; 3.0	ns	3.0	2.0; 3.5	0.005	2.5	2.0; 2.5	<0.001	2.0	1.5; 3.0	<0.001
PI-1	2.0	1.5; 2.5	ns	2.0	1.5; 2.5	ns	2.0	1.5; 2.0	0.002	2.0	1.5; 2.5	<0.001
PI-2	1.5	1.0; 1.5	ns	1.5	1.5; 2.0	0.004	1.5	1.0; 1.5	ns	1.5	1.0; 2.0	0.001
PI-3	1.5	1.0; 2.0	0.046	2.0	2.0; 2.5	ns	2.0	1.5; 2.0	0.001	2.0	1.5; 2.0	<0.001
PI-4	2.0	1.0; 2.0	ns	2.0	1.5; 2.5	0.046	2.0	1.5; 2.0	0.002	2.0	1.5; 2.0	<0.001
PI-5	1.5	1.5; 2.0	0.042	2.0	1.5; 2.5	ns	1.5	1.5; 2.0	0.04	1.5	1.5; 2.0	0.008
MPQ-A	1.0	1.0; 1.5	0.01	2.0	1.5; 2.5	0.04	1.5	1.0; 2.0	ns	1.5	1.0; 2.0	0.002
MPQ-S	1.5	1.0; 1.5	0.02	1.5	1.5; 2.0	ns	1.5	1.0; 1.5	ns	1.5	1.0; 1.5	0.003

an audiotope and expect him to benefit from listening to the tape [58]. Rather, structured audiotope programs on relaxation have been shown to be effective when the patient was first instructed on their use and encouraged to use them [59]. In the present study, therefore, craniofacial pain sufferers were provided with advice and instructions about the progressive muscle relaxation program to be followed. Acceptance of the program was rather uniform among patients. Similarly, a significant global benefit was shown for the whole sample. In agreement with this finding, the effectiveness of relaxation techniques in the management of TMD including headache, facial and musculoskeletal pain has been proven in several systematic reviews and meta-analyses [33,60-63]. Programs involving relaxation techniques to relieve pain may even be superior to that of occlusal splints in persons with acute or chronic myofascial or muscular TMD [62,64].

Psychological subtyping

The acknowledgement of the interaction between somatic complaints and psychosocial factors warrants a multidimensional approach to the classification of patients with craniofacial pain and dysfunctions. Two decades ago, Turk and Rudy [21] suggested an empirical taxonomy of chronic pain patients comprising three psychophysical subgroups that have been referred to as “dysfunctional”, “interpersonally distressed” and “adaptive copers”. They were replicated for low back pain, headache, and TMD [22]. Later, “repressive patients” were suggested to form a subtype of the “dysfunctional patients” [65]. Dysfunctional patients show high pain intensity, marked pain interference with occupational and social activities, affective distress and reduced physical activity, whereas “adaptive” patients judge their pain as less intense and interfering, are emotionally less distressed and report more activity and more control of their lives [18,21,22,54]. Other studies arrived at similar classifications of temporomandibular pain sufferers who were different in their illness beliefs, affective involvement, coping mechanisms and impact on private and professional life, but not in their objective physical findings [19,23,24]. Although meaningful subgroups of pain sufferers were consistently and reproducibly established, only few studies aimed to predict differential treatment outcome from this classification [25-30,66].

In our study, homogeneous patient clusters were identified on the basis of the whole body of questionnaire data assessed. In agreement with previous studies [19,21,22,26,66] three patient subtypes emerged which were significantly different for pain parameters and psychological profile. Cluster I subjects on average showed the lowest symptom burden as reflected by the Pain Inventory and Zerssen’s Complaint List. The irrational attitudes scores of these subjects are between those of the

two other clusters. The increased use of cognitive control and the low avoidance scores may be related to the lower subjective pain stress. Thus, this subgroup resembles the type of patient referred to in literature as “uncomplicated” [19], “low impact” [26] or “good pain control” [66] with low levels of pain, impairment and dysfunction.

Overall self-perceived pain relief after relaxation exercise was significantly lower in cluster I patients compared to the two other clusters. This was supported by the pre-post treatment findings for pain ratings. The low-symptom Cluster I patients showed the least pre-posttreatment differences. Perceived pain intensity significantly decreased like in the other groups, but pain duration, frequency and interference which were already low pre-treatment did not change. Significant improvements were also observed for the pain domains CDP and FTP and either dimension of the MPQ.

Cluster II comprised persons of younger age whose pain symptoms prevail in the fronto-temporo-mandibular region and were perceived as more severe and emotionally distressing than on average. This is consistent with findings indicating that the preference of MPQ pain descriptors is associated with pain chronicity [67] as well as pain type and localization [68]. In contrast to other TMD sufferers, patients with primarily myogenic facial pain and headache choose more affective descriptors and rate the pain as more intensive [24,52,67,68]. The higher psychosomatic burden reflected by the CL scores underpins the picture of subjects who display various pain locations of similar intensity and a number of accessory complaints. Irrational beliefs, especially a conspicuously negative self-appraisal were consistently more marked in cluster II as compared to the two other groups. Moreover, these persons experienced less social support and responded with increased withdrawal instead of activity. This may partly be due to increased sensory perception of primarily fronto-temporal pain. In summary, the negative self-image, the reduced interpersonal satisfaction, the psychosomatic burden and the affective pain component may point to premorbid affective lability and introversion. Thus, this subgroup is characterized by increased psychological maladaptation and corresponds well to the “chronic pain syndrome” [66] or “dysfunctional” type described in literature [19,21,22,69]. These patients showed significant general reductions of pain intensity, frequency, duration, and interference. Moreover, the temporomandibular and occipito-cervical pain domains were significantly improved as well as the affective dimension of the MPQ.

Interviewees belonging to group III made up nearly half of our sample. Their symptoms had existed for 9 years on average and, thus, about double as long as in the other groups. Pain was localized primarily in the occiput, neck and back areas and partly accompanied by

parafunctional activities and a higher burden of somatic complaints in comparison to cluster I. In contrast, the frequency of fronto-temporal headache and particularly of characteristic TMD signs was reduced. Pain intensity was nearly as high as in group II. Members of group III showed the highest degree of psychological normality as reflected by the Irrational Attitudes Questionnaire. Pain behavior resembled that of group I involving activity as the prevailing coping strategy. More social support was reported in comparison to the other groups. Thus, group III is primarily characterized by active coping with pain, representing the “adaptive” type of patients [19,21,66,69]. Alternatively, such patients may reflect a “stoic” profile which was described for persons with a long history of pain who show little psychological dysfunction relative to perceived pain intensity and pain-related interference [70,71]. Presumably, a stable primary personality and long-lasting “adaptation” to the burden of complaints may have led to successful coping in most of these individuals. In some patients, however, a mismatch between pain perception and psychological self-appraisal may result from a type of “Freudian repression” rather than adaptation. While pain intensity and somatic complaints are similar to those of “dysfunctional” patients, the mild cognitive symptoms mimic those of “adaptive copers” [65].

Posttreatment, cluster III patients revealed significantly reduced pain intensity, frequency, duration, and interference like cluster II patients, but pain descriptors in the MPQ remained unchanged. Furthermore, significant improvements were found for all pain domains except TMD which showed a significantly lower pretreatment level as compared to the other clusters.

All in all, the typological composition of our sample best resembles that reported by Suvinen et al. [19] who distinguished a highly distressed, psychosocially maladaptive group, a moderately distressed, behaviorally functional group, and a predominantly physical disorder group with an unremarkable psychosocial profile. Moreover, there were cluster-specific differences in the extent of change for general pain intensity and interference, the pain domains PFA, OCP and CDP, and the MPQ descriptor ratings. The lesser improvement of pain frequency, duration and interference observed in cluster I patients may be accounted for by lower baseline levels of these variables. However, significant effects were found for general pain intensity ratings and selected pain domains. Similarly, significant reductions in pre-to-posttreatment pain were found in the “low impact” subgroup of migraine headache sufferers [26]. Thus, any craniofacial patient subtype may benefit from a self-administered relaxation training. It may be applied even prior to in-depth diagnosis, in cases with unclear etiology, or in addition to conservative treatment. Nevertheless, the variable treatment outcomes observed support the call for tailored treatment

for psychologically defined subgroups or individual patients with chronic pain [14,56,72,73].

Conclusions

- Craniofacial pain sufferers may be classified into subtypes with different symptom-related and psychosocial characteristics (low symptom and psychosocial impact – high symptom and psychosocial impact – high symptom impact, low psychosocial impact).
- A self-administered relaxation training, which is cost-effective and time-saving, generally yields positive effects in patients with craniofacial pain.
- Different patient subtypes benefit differently from such training, which is most promising in patient with high symptom impact and moderate or high dysfunction.

Consent

Written informed consent was obtained from the patient’s guardian/parent/next of kin for the publication of this report and any accompanying images.

Competing interest

The authors declare that they have no competing interests.

Authors’ contributions

CK, PR, PP and CL contributed to the conception, design and coordination of the study. CK made substantial contributions to the acquisition of data and the preparation of the manuscript. CK drafted and wrote the manuscript. PR, PP and CL revised the manuscript. All authors read and approved the final manuscript.

Acknowledgements

The authors want to thank the Open Publication Fund of the University of Muenster for their support.

Author details

¹Department of Orthodontics, University Medical Centre of Regensburg, Franz-Josef-Strauß-Allee 11, Regensburg 93053, Germany. ²Department of Orthodontics, University Medical Centre of Muenster, Waldeyerstraße 30, Münster 48149, Germany.

Received: 5 August 2013 Accepted: 9 October 2013

Published: 20 October 2013

References

1. Sipilä K, Ylöstalo P, Joukamaa M, Knuuttila M: Comorbidity between facial pain, widespread pain, and depressive symptoms in young adults. *J Orofac Pain* 2006, **20**:24–30.
2. Von Korff M, Dworkin S, LeResche L, Kruger A: An epidemiologic comparison of pain complaints. *Pain* 1988, **32**:173–183.
3. Mongini F: Temporomandibular disorders and tension-type headache. *Curr Pain Headache Rep* 2007, **11**:465–470.
4. Türp J, Kowalski C, Stohler C: Temporomandibular disorders - Pain outside the head and face is rarely acknowledged in the chief complaint. *J Prosthet Dent* 1997, **78**:592–595.
5. Bernhardt O, Gesch D, Schwahn C, Mack F, Meyer G, John U, Kocher T: Risk factors for headache, including TMD signs and symptoms, and their impact on quality of life. Results of the Study of Health in Pomerania (SHIP). *Quintessence Int* 2005, **36**:55–64.
6. Jensen R, Stovner L: Epidemiology and comorbidity of headache. *Lancet Neurol* 2008, **7**:354–361.

7. Glaros A, Urban D, Locke J: **Headache and temporomandibular disorders: evidence for diagnostic and behavioural overlap.** *Cephalalgia* 2007, **27**:542–549.
8. Dando W, Branch M, Maye J: **Headache disability in orofacial pain patients.** *Headache* 2006, **46**:322–326.
9. Mongini F, Ciccone G, Ibertis F, Negro C: **Personality characteristics and accompanying symptoms in temporomandibular joint dysfunction, headache, and facial pain.** *J Orofac Pain* 2000, **14**:52–58.
10. Dao TT, LeResche L: **Gender differences in pain.** *J Orofac Pain* 2000, **14**:169–184.
11. Munce S, Stewart D: **Gender differences in depression and chronic pain conditions in a national epidemiologic survey.** *Psychosomatics* 2007, **48**:394–399.
12. Magnusson T, Egermark I, Carlsson GE: **A longitudinal epidemiologic study of signs and symptoms of temporomandibular disorders from 15 to 35 years of age.** *J Orofac Pain* 2000, **14**:310–319.
13. John MT, Reissmann DR, Schierz O, Wassell RW: **Oral health-related quality of life in patients with temporomandibular disorders.** *J Orofac Pain* 2007, **21**:46–54.
14. Dworkin SF, Huggins KH, Wilson L, Mancl L, Turner J, Massoth D, LeResche L, Truelove E: **A randomized clinical trial using research diagnostic criteria for temporomandibular disorders-axis II to target clinic cases for a tailored self-care TMD treatment program.** *J Orofac Pain* 2002, **16**:48–63.
15. Forssell H, Kalso E: **Application of principles of evidence-based medicine to occlusal treatment for temporomandibular disorders: are there lessons to be learned?** *J Orofac Pain* 2004, **18**:9–22.
16. Von Korff M: **Health services research and temporomandibular pain.** In *Temporomandibular disorders and related pain conditions*. Edited by Sessle BJ, Bryant PS, Dionne RA. Seattle: IASP Press; 1995:227–236.
17. Dworkin SF: **Psychosocial issues.** In *From basic science to clinical management*. Edited by Lavigne G, Sessle BJ, Dubner R. Carol Stream, Ill: Quintessence; 2001:115–128.
18. Oakley ME, McCreary CP, Clark GT, Holston S, Glover D, Kashima K: **A cognitive-behavioral approach to temporomandibular dysfunction treatment failures: a controlled comparison.** *J Orofac Pain* 1994, **8**:397–401.
19. Suvinen TI, Hanes KR, Gerschman JA, Reade PC: **Psychophysical subtypes of temporomandibular disorders.** *J Orofac Pain* 1997, **11**:200–205.
20. Sessle BJ: **Integration of basic sciences into the predoctoral curriculum to study temporomandibular disorders and orofacial pain.** *J Orofac Pain* 2002, **16**:181–184.
21. Turk DC, Rudy T: **Toward an empirically derived taxonomy of chronic pain patients: integration of psychological assessment data.** *J Consult Clin Psychol* 1988, **56**:233–238.
22. Jamison R, Rudy T, Penzien D, Mosley T: **Cognitive-behavioral classifications of chronic pain: replication and extension of empirically derived patient profiles.** *Pain* 1994, **57**:277–292.
23. Schulte JK, Anderson GC, Hathaway KM, Will TE: **Psychometric profiles and related pain characteristics of temporomandibular disorder patients.** *J Orofac Pain* 1993, **7**:247–253.
24. Mongini F, Italiano M, Raviola F, Mossolov A: **The McGill Pain Questionnaire in patients with TMJ pain and with facial pain as a somatoform disorder.** *Cranio* 2000, **18**:249–256.
25. Sanders S, Brena S: **Empirically derived chronic pain patient subgroups: the utility of multidimensional clustering to identify differential treatment effects.** *Pain* 1993, **54**:51–56.
26. Davis P, Reeves J, Graff-Radford S, Hastie B, Naliboff B: **Multidimensional subgroups in migraine: differential treatment outcome to a pain medicine program.** *Pain Med* 2003, **4**:215–222.
27. Swimmer G, Robinson M, Geisser M: **Relationship of MMPI cluster type, pain coping strategy, and treatment outcome.** *Clin J Pain* 1992, **8**:131–137.
28. Gatchel R, Noe C, Pulliam C, Robbins H, Deschner M, Gajraj NM, Vakharia AS: **A preliminary study of multidimensional pain inventory profile differences in predicting treatment outcome in a heterogeneous cohort of patients with chronic pain.** *Clin J Pain* 2002, **18**:139–143.
29. Denison E, Asenlöf P, Sandborgh M, Lindberg P: **Musculoskeletal pain in primary health care: subgroups based on pain intensity, disability, self-efficacy, and fear-avoidance variables.** *J Pain* 2007, **8**:67–74.
30. Riley JL, Robinson M, Geisser M, Wittmer V, Smith A: **Relationship between MMPI-2 cluster profiles and surgical outcome in low-back pain patients.** *J Spinal Disord* 1995, **8**:213–219.
31. DeBar LL, Vuckovic N, Schneider J, Ritenbaugh C: **Use of complementary and alternative medicine for temporomandibular disorders.** *J Orofac Pain* 2003, **17**:224–236.
32. Raphael KG, Klausner JJ, Nayak S, Marbach JJ: **Complementary and alternative therapy use by patients with myofascial temporomandibular disorders.** *J Orofac Pain* 2003, **17**:36–41.
33. Orlando B, Manfredini D, Salvetti G, Bosco M: **Evaluation of the effectiveness of biobehavioral therapy in the treatment of temporomandibular disorders: a literature review.** *Behav Med* 2007, **33**:101–118.
34. Türp JC, Kowalski CJ, Stohler CS: **Treatment-seeking patterns of facial pain patients: many possibilities, limited satisfaction.** *J Orofac Pain* 1998, **12**:61–66.
35. Carlson CR, Okeson JP, Falace DA, Nitz AJ, Curran SL, Anderson D: **Comparison of psychologic and physiologic functioning between patients with masticatory muscle pain and matched controls.** *J Orofac Pain* 1993, **7**:15–22.
36. Linton SJ: **An overview of psychosocial and behavior factors in neck and shoulder pain.** *Scand J Rehabil Med* 1995, **32**(Suppl):67–78.
37. Aaron LA, Turner JA, Mancl LA, Sawchuk CN, Huggins KH, Truelove EL: **Daily pain coping among patients with chronic temporomandibular disorder pain: an electronic diary study.** *J Orofac Pain* 2006, **20**:125–137.
38. Riley JL, Myers CD, Currie TP, Mayoral O, Harris RG, Fisher JA, Gremillion HA, Robinson ME: **Self-care behaviors associated with myofascial temporomandibular disorder pain.** *J Orofac Pain* 2007, **21**:194–202.
39. Geissner E: **The pain perception Scale - a differentiated and change-sensitive scale for assessing chronic and acute pain.** *Rehabilitation* 1995, **34**:35–43.
40. von Zerssen D: **Clinical Self-Rating Scales (CSRS) of the Munich Psychiatric Information System (PSYCHIS München).** In *Assessment of depression*. Edited by Sartorius N, Ban T. Berlin-New York: Springer; 1986:270–303.
41. Klages U: **Fragebogen irrationaler Einstellungen (FIE), Irrational Attitudes Questionnaire.** Goettingen, Germany: Hogrefe; 1989.
42. Klages U: **Fragebogen zum Schmerzverhalten (FSV), Pain Behavior Questionnaire.** *Diagnostica* 1989, **35**:351–353.
43. John MT, Hirsch C, Reiber T, Dworkin S: **Translating the research diagnostic criteria for temporomandibular disorders into German: evaluation of content and process.** *J Orofac Pain* 2006, **20**:43–52.
44. Bellach BM: **Der Bundes-Gesundheitssurvey von 1998. Erfahrungen, Ergebnisse und Perspektiven.** *Gesundheitswesen* 1999, **61**:55–56. Spec No.
45. Johansson A, Unell L, Carlsson GE, Soderfeldt B, Halling A: **Gender difference in symptoms related to temporomandibular disorders in a population of 50-year-old subjects.** *J Orofac Pain* 2003, **17**:29–35.
46. Kuttilla M, Niemi P, Kuttilla S, Alanen P, LeBell Y: **TMD treatment need in relation to age, gender, stress, and diagnostic subgroup.** *J Orofac Pain* 1998, **12**:67–74.
47. John M, Miglioretti D, LeResche L, Von Korff M, Critchlow C: **Widespread pain as a risk factor for dysfunctional temporomandibular disorder pain.** *Pain* 2003, **102**:257–263.
48. Von Korff M, Ormel J, Keefe F, Dworkin S: **Grading the severity of chronic pain.** *Pain* 1992, **50**:133–149.
49. Türp J, Kowalski C, Stohler C: **Generic pain intensity scores are affected by painful comorbidity.** *J Orofac Pain* 2000, **14**:47–51.
50. Turner JA, Brister H, Huggins K, Mancl L, Aaron L, Truelove E: **Catastrophizing is associated with clinical examination findings, activity interference, and health care use among patients with temporomandibular disorders.** *J Orofac Pain* 2005, **19**:291–300.
51. Turner JA, Holtzman S, Mancl L: **Mediators, moderators, and predictors of the therapeutic change in cognitive-behavioral therapy for chronic pain.** *Pain* 2007, **127**:276–286.
52. Jerjes W, Madland G, Feinmann C, Hopper C, Kumar M, Upile T, Kudari M, Newman S: **A psychological comparison of temporomandibular disorder and chronic daily headache: are there targets for therapeutic interventions?** *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 2007, **103**:367–373.
53. Schwartz SM, Gramling SE: **Cognitive factors associated with facial pain.** *Cranio* 1997, **15**:261–266.
54. Epker J, Gatchel RJ: **Coping profile differences in the biopsychosocial functioning of patients with temporomandibular disorder.** *Psychosom Med* 2000, **62**:69–75.
55. Michelotti A, Steenks MH, Farella M, Parisini F, Cimino R, Martina R: **The additional value of a home physical therapy regimen versus patient education only for the treatment of myofascial pain of the jaw muscles: short-term results of a randomized clinical trial.** *J Orofac Pain* 2004, **18**:114–125.

56. Michelotti A, de Wijer A, Steenks M, Farella M: **Home-exercise regimes for the management of non-specific temporomandibular disorders.** *J Oral Rehabil* 2005, **32**:779–785.
57. Mulet M, Decker KL, Look JO, Lenton PA, Schiffman EL: **A randomized clinical trial assessing the efficacy of adding 6 × 6 exercises to self-care for the treatment of masticatory myofascial pain.** *J Orofac Pain* 2007, **21**:318–328.
58. Okeson JP, Moody PM, Kemper JT, Haley JV: **Evaluation of occlusal splint therapy and relaxation procedures in patients with temporomandibular disorders.** *J Am Dent Assoc* 1983, **107**:420–424.
59. Carrington P, Collings GH, Benson H, Robinson H, Wood LW, Lehrer P, Woolfolk RL, Cole JW: **The use of meditation - relaxation techniques for the management of stress in a working population.** *J Occup Med* 1980, **22**:221–231.
60. Türp J, Jokstad A, Motschall E, Schindler H, Windecker-Gétaz I, Ettlin D: **Is there a superiority of multimodal as opposed to simple therapy in patients with temporomandibular disorders? A qualitative systematic review of the literature.** *Clin Oral Implants Res* 2007, **18**(Suppl 3):138–150.
61. Myers CD: **Complementary and alternative medicine for persistent facial pain.** *Dent Clin North Am* 2007, **51**:263–274.
62. Medicott MS, Harris SR: **A systematic review of the effectiveness of exercise, manual therapy, electrotherapy, relaxation training, and biofeedback in the management of temporomandibular disorder.** *Phys Ther* 2006, **86**:955–973.
63. Sierpina V, Astin J, Giordano J: **Mind-body therapies for headache.** *Am Fam Physician* 2007, **76**:1518–1522.
64. Brooke RI, Stenn PG: **Myofascial pain dysfunction syndrome: how effective is biofeedback-assisted relaxation training?** In *Advances in pain research and therapy*, Volume 5. Edited by Bonica JJ, Lindblom U, Iggo A. New York: Raven; 1983:809–812.
65. Burns J, Kubilus A, Bruehl S, Harden R: **A fourth empirically derived cluster of chronic pain patients based on the multidimensional pain inventory: evidence for repression within the dysfunctional group.** *J Consult Clin Psychol* 2001, **69**:663–673.
66. Klapow J, Slater M, Patterson T, Doctor J, Atkinson J, Garfin S: **An empirical evaluation of multidimensional clinical outcome in chronic low back pain patients.** *Pain* 1993, **55**:107–118.
67. Türp JC, Kowalski CJ, Stohler CS: **Pain descriptors characteristic of persistent facial pain.** *J Orofac Pain* 1997, **11**:285–290.
68. Mongini F, Italiano M: **TMJ disorders and myogenic facial pain: a discriminative analysis using the McGill Pain Questionnaire.** *Pain* 2001, **91**:323–330.
69. Suvinen TI, Reade PC, Hanes KR, Könönen M, Kempainen P: **Temporomandibular disorder subtypes according to self-reported physical and psychosocial variables in female patients: a re-evaluation.** *J Oral Rehabil* 2005, **32**:166–173.
70. Cook A, Degood D: **The cognitive risk profile for pain: development of a self-report inventory for identifying beliefs and attitudes that interfere with pain management.** *Clin J Pain* 2006, **22**:332–345.
71. Sheffer CE, Deisinger JA, Cassisi JE, Lofland K: **A revised taxonomy of patients with chronic pain.** *Pain Med* 2007, **8**:312–325.
72. Turk D, Rudy T, Kubinski J, Zaki H, Greco C: **Dysfunctional patients with temporomandibular disorders: evaluating the efficacy of a tailored treatment protocol.** *J Consult Clin Psychol* 1996, **64**:139–146.
73. Turk DC, Swanson K, Tunks E: **Psychological approaches in the treatment of chronic pain patients - when pills, scalpels, and needles are not enough.** *Can J Psychiatr* 2008, **53**:213–223.

doi:10.1186/1746-160X-9-31

Cite this article as: Kirschneck *et al.*: Psychological profile and self-administered relaxation in patients with craniofacial pain: a prospective in-office study. *Head & Face Medicine* 2013 9:31.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

